



## **Implant Post-Operative Instructions**

A dental implant has been placed in your mouth. The Implant may be located above or below the tissue. This type of implant has been selected for your situation because of the bone available to place an implant. The implant will usually take a period of one to four months to heal; depending upon your body's healing properties and the type of implant surgery.

If IV sedation was used for your procedure, you cannot drive a car for the remainder of the day.

If sutures were placed around your implant, they will be removed and checked by your doctor in one to two weeks.

Usually, you are able to wear your present partial or flipper, if applicable. Sometimes it is necessary to leave it in the night of the surgery. It is important to keep the appliance as clean as possible during the healing period. Please ask the doctor if you have any questions.

Every consideration must be given to keep the surgical site clean and free of food particles.

## **Instructions for oral care and other helpful information:**

**NO SMOKING!** Smoking is to be avoided for the time specified by the doctor. Smoking increases the heat in the surgical site and significantly lowers the body's ability to heal.

**AVOID** these after surgery: Alcohol with post-operative medications, commercial mouth rinses and very hot fluids.

Gentle rinsing of the mouth should be started the day after surgery. Frequent gentle rinsing with lukewarm salt water will aid the healing process (add one half teaspoon of salt to a 6 oz glass of water). Avoid the use of a water-pik tooth brush.

**Pain:** A certain amount of pain must be expected with all types of surgery. An appropriate medication has been prescribed for you. Please take it according to the directions. It is advised that you do not drive while take the pain prescriptions. However, if only Ibuprofen (Motrin) or Tylenol is needed, most people are able to drive without a problem.

**Swelling:** Some swelling and minimal bruising is possible and is to be expected. IT IS NOT UNUSUAL. In most cases, swelling can be prevented/controlled. Apply the ice pack that has been given to you for a period of 20 minutes on and 20 minutes off during the day for the next two days. The application of ice to the outside of the face over the surgical area will minimize swelling. If after five days you still have unusual swelling or pain, please call the office.

**Bleeding:** A small amount of bleeding following surgery should not alarm you. If the bleeding is excessive or continuous, please call us at once. Vigorous rinsing of the mouth prolongs bleeding by removing the clotting blood, so when rinsing you mouth, do it gently.

**NO** drinking through straws. The use of a straw creates negative pressure in your mouth and will tend to loosen the sutures. Also the suctioning action can cause you to suck the blood clot out and cause a dry socket.

**IF YOU HAVE QUESTIONS AT ANY TIME, PLEASE CONTACT OUR OFFICE AT (517) 748-9440**

**Diet:** Following surgery it is best to restrict your diet to fluids and soft foods for the first day. Normal diet may then be resumed the following day, but you will want to avoid chewing on the implant site until the tissue is completely healed. Soft foods such as Jell-O, pudding, mashed potatoes, scrambled eggs and soups are suggested. If you have difficulty chewing, try blenderized foods or diet supplements such as Carnation instant breakfast and ensure.

**Implant Follow-up:** We will see you for the periodic check-ups until the implant has healed. This is usually for a period of one to four months, depending on your implant surgery.